PLANTING DATES FOR CLOVER, ALFA-RACK PLUS, EXTREME, NO-PLOW, FUSION, CHIC MAGNET and EDGE

1 Call for planting dates

2 Apr 1 - July 1

3 Apr 15 - June 15 Aug 1 - Sept 1

4 Coastal: Feb 1 - Mar 15 Sept 1 - Oct 15 Southern Piedmont: Feb 15 - Apr 1 Aug 15 - Oct 1 Mountain Valleys: Mar 1 - Apr 15 Aug 1 - Sept 15

5 Feb 1 - Apr 1 Aug 1 - Sept 30 6 Feb 1 - Apr 15 Sept 1 - Nov 1

 North: Mar 15 - May 1 Aug 1 - Sept 15
 South: Mar 1 - Apr 15 Aug 15 - Oct 15 20

19

2

- 8 Apr 1 June 15 July 15 - Sept 5
- 9 Apr 1 May 15 Aug 1 -Sept 15
- 10 Mar 20 May 15 Aug 1 - Sept 15
- 1 Sept 15 Nov 15

Feb 5 - Mar 1
 North: Sept 5 - Nov 15
 South: Sept 25 - Nov 15
 Feb 15 - Apr 1

2

13

6

14

- Sept 1 Oct 30 North: Sept 15 - Nov 15 South: Sept 25 - Nov 15
- Feb 1 Mar 1
 Coastal: Sept 25 -Oct 15
 Piedmont: Sept 1 -Oct 5
 Mountain Valleys: Aug 25 - Oct 15
- North: Sept 25 Nov 25 South: Oct 5 - Nov 30
 Mar 1 - May 15 Aug 1 - Sept 15
 Feb 1 - Apr 15 Aug 20 - Sept 30
 Apr 15 - June 15 July 1 - Aug 15
 May 15 - July 1
 May 1 - June 15 July 1 - Aug 15
 May 15 - July 1

22

21

4

15

10

18

12

PLANTING DATES FOR DESTINATION, VISION, PURE ATTRACTION, SECRET SPOT, BOWSTAND and WINTER PEAS



16

Aug 15 - Nov 1
North: Aug 1 - Sept 30 South: Aug 15 - Oct 15
July 15 - Sept 5
Aug 1 - Sept 15
Aug 1 - Sept 15

- 1 Sept 15 Nov 15
 - North: Sept 5 Nov 15 South: Sept 25 - Nov 15



22 July 1 - Aug 15

PLANTING DATES FOR WINTER-GREENS, TALL TINE TUBERS, RAVISH RADISH and BEETS & GREENS





PLANTING DATES FOR IMPERIAL POWERPLANT, TURKEY SELECT, SUNN HEMP AND CONCEAL



*Do not plant PowerPlant, Sunn Hemp or Conceal until soil temperatures reach a constant 65 degrees F. Wait as long as necessary for soil temperatures to reach a constant 65 degrees F before planting.

PLANTING DATES FOR IMPERIAL WHITETAIL OATS PLUS

Use the map below as a guideline for when to plant Imperial Whitetail Oats Plus in your area. For best results, wait to plant until excessively hot, droughty summer weather has passed.

Imperial Whitetail Oats Plus is highly cold-tolerant and designed to provide abundant forage from fall into spring in the southern U.S. and from fall into winter in colder climates.

